

The effect of watery extract *Cuminum* on the testes of the local Rabbits (*Oryctolagus cuniculus*)

Hudayiaat Abd al Razzaq¹, Walaa Qassem Taemeh², Walaa Fadil Obead³
University of Kerbala, College of Veterinary Medicine, Department of Anatomy

Abstract — This study was conducted to investigate the effect of watery extract of Cumin plant (200 mg/kg B.W.) on some histological changes of the male reproductive system in albino rabbits. Sixteen adult male rabbits weighting about (200-250 g) and 12-15 weeks age were used in the present study and divided into (2) groups, 8 rabbits per group, group1: serve as a control group and given normal diet with drinking water, group2: Given Cumin extract (200 mg/kg B.W.) through oral tubular feeding, the experiment were continued 60 days for all groups. The results of the current study showed significant decrease ($P<0.05$) in serum testosterone compared with control group. As for ectopic tissue, water extract (200 mg/kg) has resulted in satisfactory tissue changes of the testes. The water extract of Cumin plant acts as an antioxidant through its important role in curbing the harmful effects of some types of free radicals within the body and histological changes and functions of the male reproductive system of rabbits. The results of histological diagnosis of testes in male rabbits that giving cumin for long periods leads to a decrease in the level of testosterone

Keywords — Testis, effect of *Cuminum*, Rabbits.

I. INTRODUCTION

CUMIN is a multi-purpose aromatic plant used worldwide as a food additive and represents the popularity of spices to flavor different preparations, including cheese, pickles, soups, beans, and alcoholic beverages. It is also used in traditional medicine, especially in veterinary medicine [1]. Cumin seeds have shown anti-oxidant properties, as well as useful in indigestion, diarrhea and cholera, as a treatment against indigestion and colic [2], also Cumin seed oil used as active oil against microbes [3]. These medicinal benefits were generally referenced to their content of active ingredients and strong action such as terpenes, phenols, and flavonoids, it also contains essential fatty acid and various classes of compounds, including terpenes, alcohol, phenols and aldehydes more precisely known as aldehyde Cumin: cumin aldehyde, eugenol, B-Benin and some other minor compounds [4]. The botanical kingdom is the actual source of most of the drugs and other active pharmaceutical molecules that are waiting to be discovered. Over the past decade, the therapeutic use of herbal medicines has gained momentum worldwide due to the high toxicity of many chemically manufactured pharmaceuticals, resulting in a sudden increase in the number of herbal medicine manufacturers [5]. The cumin product has demonstrated an active antioxidant capacity, with the ability to suppress free radicals and fatty peroxides [6]. It has the ability

to reduce cholesterol levels in plasma for diabetic rats [7]. *Cuminum cyminum* was used as an antioxidant in male rabbits by measuring the level of testosterone [8] mentioned that the testis of adult rams is exist within a specialized pouch of skin, the scrotum. The testes are enclosed by capsule, the tunica albuginea which is composed of dense collagenous tissue. This capsule is covered by, tunica serosa, the connective tissue of which blends with the tunica albuginea. The tunica albuginea is continuous with the loose collagenous tissue of mediastinum testis and dividing the testis into the lobules (lobuli testis). The tubules of the lobule (seminiferous tubules and rete testis) are surrounded by loose collagenous tissue with many reticular fibers [9] and [10] reported that the testicular seminal pathway in camel is divided into two portions; an intratesticular and extra testicular portion. The extra testicular portion consist of the rete testis, an extension of the intratesticular portion and an elongated sac from which 6-7 ductuli efferent originated. Studies which are done by [11] revealed that the long straight seminiferous tubules in the male camel are lined with simple cuboidal cells to simple columnar. The different types of germ cells of the sperm lie in seminiferous tubules and represents different phases within the sperm development. The wide base of Sertoli cells stands on the basement membrane of seminiferous tubules and it is usually that the surface of Sertoli cell contains the old spermatids. The interstitial cells (Leydig cells) are being polygonal, organized in groups or nest and well supplied by blood capillaries and occupies the spaces that lies among the seminiferous tubules. Those researchers also noticed that the rete testis is divide in to a part inside the testis and external part. The external part of the rete testis forms a sac in the meeting of the ductuli efferences. Spermatocytogenesis begins with spermatogonia on the basement membrane of seminiferous tubules and proceed toward the lumen. The formation of spermatid marks the end of spermatogenesis and beginning of the spermiogenesis [11].

II. MATERIALS AND METHODS

A. Animals Housing and Experimental Design

sixteen adult male albino rabbits weighting about (200-250 g) were used in the present study and divided into (2) groups, 8 rabbits per group, group1: serve as a control group and given normal diet with drinking water, group2: given Cumin extract (200 mg/kg B.W.) through oral tubular feeding and the experiment were continued 60 days for all groups.

Collection of Plant and Preparation of

Extract:

Cumin powders were obtained from the local market in Kerala city. The water extract of Cumin powder was prepared based on [14].

Blood Sampling:

At the end of each experiment, the animals were anesthetized with ketamine and xylazine for several minutes, after that the front and hind limbs of the animal were fixed with pins. blood samples were taken directly from the heart by cardiac stenosis, collected about 8-10 ml of blood, placed in Test tubes free of anticoagulants left for about a quarter of an hour at room temperature [15] In the centrifuge at 3000 / rpm for 15 minutes and take serum, and kept at (-20 °C) in new, clean plastic tubes (Plane Tubes) until required biochemical tests were carried out.

Dissection and Removal of Organs:

Testis was pulled using special forceps and removed with scissors at the level of the groin duct, and then placed in a Petri dish containing the saline salt solution (0.9% NaCl) to separate it from the surrounding grease, and were kept in formalin at a concentration of 10% for the conduct of tissue study.

Histological Sections Preparation:

Samples of testes were directly fixed in 10% buffered formalin for 24 hours and then processed for paraffin method by dehydrating through ascending concentrations of ethanol (60%, 70%, 80%, 90%, 95% and 100%), cleared in xylene, infiltrated in paraffin wax and finally embedded in paraffin wax. Sections were cut at 5µm thickness with a rotary microtome (Hunting Don, Bright.UK). The sections were stained by hematoxylin and eosin (H&E) method [16].

Statistical analysis:

The results were statistically analyzed using the Statistical Analysis Package for Social Science (SPSS). In order to extract the differences between the experimental groups with emphasis on these differences by extracting the standard error (Stander Error) SE Statistical analyzes were conducted according to Duncan and others [17].

III. RESULTS AND DISCUSSION

The choose of rabbit in this research is considered as an important animal for an anatomical representative of class Mammalia in comparative anatomy classes and its size makes it a convenient research animal. Each testis is found in a large sac or scrotum just ventral to the anus (fig 1,2). The scrotum when cut shows the following layers from the exterior to the inner

wall: (1) Skin, (2) Cremaster muscle and fascia, and (3) Tunica vaginalis, a thick portion of the peritoneum that lines the wall of the scrotum or parietal lamina and covers the testes or visceral lamina. This membrane is continuous with the serous membranes of the peritoneal cavity (fig 1,2) A thick fibrous connective tissue capsule (fig 3), the tunica albuginea,

covers each testis. The inner part of this capsule is the tunica

vascular, a loose connective tissue layer that contains blood vessels. Each testis is divided into lobules by incomplete connective tissue septa and there is frequently intercommunication between the lobules. Each lobule is occupied by seminiferous tubules (fig 3) enmeshed in a web of loose connective tissue that is rich in blood and lymphatic vessels, nerves and interstitial or Leydig cells. All phases of event cycle of spermatogenesis and spermiogenesis in rabbit are observed within the seminiferous tubules. The seminiferous epithelium or germ cells show a regular organization, with the basally located spermatogonia, together with spermatocytes, occupying half of the layer of the spermatogenic epithelium. A great number of spermatids and spermatozoa are embedded in the cytoplasmic prolongations of the Sertoli cells. Early spermatids possess a central, rounded nucleus, with granular chromatin. The differentiation of the spermatids into spermatozoa involves the events of nuclear elongation, formation of the acrosomal and axonemal complexes and elimination of residual cytoplasm.

Histological effect of treatment on testis:

The normal feature of the testis in the control group were showed in the figure (3,4,5,6,7,8) in which healthy histological structure of rabbit testis having a germinal epithelium undergoing cell division and well-formed spermatids are seen. While microscopic examination of the testis tissues of the cumin group observed many changes included degenerative in some spermatogenic cells, large number of cellular debris was collected in the tubular cavity, seminiferous tubules showed few spermatozoa in the lumens, detaching of the spermatogonia from the basal lamina in

some places, an increase in the space between the seminiferous tubules, hemorrhage and decreasing in the number of Leydig cells between them, highly vacuolated spermatogonia also were seen (fig. 9,10,11,12).

These results were disagreed with [18]. The appearance of degeneration and necrosis was explained by the inhibition of cumin for mitochondrial action by disruption of energy production and thus the effect on the sodium pump and the effect of the plasma membrane or the decrease in the pH of the cells through the increase in the production of lactic acid, produces proteins and thus inhibits the formation of the plasma membrane and the occurrence of necrosis. and decreasing in the number of Leydig cells between them, highly vacuolated spermatogonia also were seen (fig. 9,10,11,12). These results were disagreed with [18]. The appearance of degeneration and necrosis was explained by the inhibition of cumin for mitochondrial action by disruption of energy production and thus the effect on the sodium pump and the effect of the plasma membrane or the decrease in the pH of the cells through the increase in the production of lactic acid, produces proteins and thus inhibits the formation of the plasma membrane and the occurrence of necrosis.

Effect of treatment on Testosterone hormone:

A significant decrease in the level of testosterone in serum of male rabbits was observed in treatment group compared to the significant events ($P < 0.05$) in serum compared to control

group.

The results of this study showed a significant decrease ($P < 0.05$) in the serum testosterone concentration of male rabbits in the treatment group compared with control group. this result is inconsistent with the result of [18]. Decrease in the concentration of sex hormones may be due to the rise of active oxygen classes in the Leydig cells, which inhibit the cloning of the steroidogenic acute protein in the mitochondrial membranes in the Leydig cells which plays an important role in the process of hormone synthesis within the Leydig cells [19]. There is a significant decrease in the treatment group with cummin, the total number of sperm compared with the control. which is inconsistent with the results of the researchers [18]. The effective oxygen classes break down the Leydig cells responsible for the secretion of testosterone in the testis, which leads to a reduction in the level of hormone responsible for the functions of the prostate and testes, decrease in the number of sperm as well as the destruction of cells lining the spermatozoa (Sertoli cells and sperm generated cells) and thus impede sperm maturity [20].




Figure 1: show the significant differences ($P < 0.05$) between groups in The Testosterone Hormone serum level




Figure 2: shows the testis of the rabbit




Figure 3: dissecting rabbit shows: a-right testis, b-left testis, c- penis, d-anal gland




Figure 4: This figure shows a-capsule, b-seminiferous tubules, c- connective tissue septa, d- blood artery, H&E stain, 100x.




Figure 5: This figure show a-A-Sertoli cell, b-B-spermatogonia H&E stain 400x




Figure 6: This figure shows a-Leydig cell H&E stain 400x




Figure 7: This figure shows a-seminiferous tubules, b-spermatid H&E stain 100x




Figure 8: This figure shows a-epididymus, b-sperm H&E stain 100x




Figure 9: This figure shows a-sperm H&E stain 400x

IV. DISCUSSION

- 1- The Cumin are harmful on the male genital system.
- 2- The Cumin are harmful on the testosterone level hormone.

There is a need for further proposing studies which may include:

- 1- Study the effects of cumin in female genital system in different animals.
- 2- Study the effects of cumin in male genital system with stress in different animals.

REFERENCES

- [1] Sultana, S.; Ripa, F. A.; Hamid, K. (2010). Comparative antioxidant activity study of some commonly used spices in Bangladesh. *Pak. J. Biol. Sci.*, 13(7), 340.
- [2] . El-Ghorab, A. H.; Nauman, M.; Anjum, F. M.; Hussain, S.; Nadeem, M. (2010). A comparative study on chemical composition and antioxidant activity of ginger (*Zingiberofficinale*) and cumin (*Cuminum cyminum*). *J. agricul. food chem.*, 58(14), 8231-8237.
- [3] Abbaszadegan, A.; Gholami, A.; Ghahramani, Y.; Ghareghan, R.; Ghareghan, M.; Kazemi, A.; Ghasemi, Y. (2016). Antimicrobial and cytotoxic activity of *Cuminum cyminum* as an intracanal medicament compared to chlorhexidine gel. *Ira.endod. J.*, 11(1), 44.
- [4] Oroojalian, F.; Kasra-Kermanshahi, R.; Azizi, M.; Bassami, M. R. (2010). Phytochemical composition of the essential oils from three Apiaceae species and their antibacterial effects on food-borne pathogens. *Food chem.*, 120(3), 765-770.

- [5] Nasreen, S. and Radha, R. (2011). *Assessment of Quality of Withaniasomnifera Dunal (Solanaceae) Pharmacognostical and physicochemical profile. Int. J. Phar. Pharm. Sci., 3(2), 152-155.*
- [6] Deepak. (2013). *Importance of Cuminum cyminum L. and Carumcarvi L. in traditional medicaments-a review. Indian J. Trad. Knowl., 12(2), 300-307.*
- [7] Dhandapani, S.; Subramanian, V. R.; Rajagopal, S.; Namasivayam, N. (2002). *Hypolipidemic effect of Cuminum cyminum L. on alloxan-induced diabetic rats. Pharmacol. Res., 46(3), 251-255.*
- [8] Obead.W.F. (2009). *Study of seminiferous epithelium and spermatogenesis in adult male rabbit. Msc. a thesis submitted to the college of veterinary medicine university of Baghdad, Baghdad-Iraq.*
- [9] Alberts B, Johnson, A. Lewis, J. Raff, M. Roberts, K. and Watter, P. (2002). *Molecular Biology of the Cell. 4th, ed. New York; Garland Science. 1065-1126.*
- [10] Wing-Yee, Lui., Dolores, D., Mruk, Will, M. Lee. and C. Yan Cheng (2003). *Adherens Junction Dynamics in the Testis and Spermatogenesis. J. of Andr, Vol. 24, No. 1. 25*
- [11] Al-Samarrae, N.S. and Sawad. (1996). *Some anatomical and histological studies on the testes of the camel (Camelus dromedarius). Iraqi Journal of Veterinary Sciences., 9: 73-82.*
- [12] V. Venkatesh, J. D Sharma and Raka Kamal (2002). *A Comparative Study of Effect of Alcoholic Extracts of Sapindus emarginatus, Terminalia belerica, Cuminum cyminum and Allium cepa on Reproductive Organs of Male Albino Rats. Asian J. Exp. Sci, Vol. 16, No. 1&2, 51-63.*
- [13] Beena Khillare and Ashish Ranjan Singh (2014). *Spermicidal activity of elettaria cardamomum and cuminum cyminum seed extracts and assessment of sperm function in albino rats International Journal of Pharmacognosy Vol. 1(4): 258-65.*
- [14] Anesini, C. and Perez, C. (1993). *Screening of plant used in argentine folk medicine for antimicrobial activity. J. Ethnopharmacol., 39(2): 119-128.*
- [15] Bristow, A. F. (1998). *Standardization of protein hormone immunoassays: current controversies. proceeding of the UKNEQAS Endocrinology meeting ACB , 3: 66-73*
- [16] Bancroft, J. D. and Gamble, M. (2008). *Theory and practice of histological techniques. Elsevier Health Sciences. 16(2): 16-10.*
- [17] Steel, R.G.D. and Torries, J.H. (1980). *Principle and Procedures of Statistics: A Biochemical Approach. 2nd edition, McGraw-Hill Book Company Inc., New York, USA.*
- [18] Othman, A.S. Alfahdawi.; Muneef, S.A. Aljanabi.; Treefa F. I. (2018). *Physiological and histological study of the effect of Cumin Cyminum watery extract and vitamin E on the male reproductive system in rats exposed to oxidative stress. Tikrit Journal of Pure Science 23(3)*
- [19] Sakhaee, E.; Emadi, L.; Azari, O.; Kheirandish, R.; Esmaili Nejad, M. R.; ShafieiBafti, H. (2016). *Effects of Cuminum cyminum L. essential oil on some epididymal sperm parameters and histopathology of testes following experimentally induced copper poisoning in mice. Andrologia, 48(5), 542-547.*
- [20] Gupta, R. S.; Saxena, P.; Gupta, R.; Kachhawa, J. B. (2011). *Evaluation of reversible contraceptive activities of Cuminumcyminum in male albino rats. Contraception, 84(1), 98-107.*